

The porous space structure of domanik shales in the east of Russian plate

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© SGEM2018. Because of the hydrocarbon reserves reducing from traditional reservoirs in Russia, much attention is paid now to unconventional resources, such as the Domanik oil shales of the Volga-Ural Basin. The Domanik deposits consist of siliceous-clay-carbonate rocks enriched with organic matter and other components of oil in the dispersed state. They are Middle Frasnian source rocks and have low porosity and permeability. Domanik shales contains many oil occurrences and rare fields, which is primarily explained by the absence of reservoirs. The increasing significance of shale oil and gas has led to the need for deeper understanding of Domanik porous space structure. The porous space of 5 samples from Domanik shales was researched using microCT and SEM. The absolute porosity of shale samples varies from 0.66 to 1.87. The pore size distribution differs depending on the composition of the rock: carbonate rich samples have less pores of 0.01 mm than organic rich ones. The pore size of the organic matter depends on the level of maturation. The authors propose a classification of pore space structure for the Domanik shales. The most important type for hydrocarbon exploration are the leaching cracks which related to carbonate rich shales.

<http://dx.doi.org/10.5593/sgem2018/1.4/S06.118>

Keywords

Domanik, MicroCT, Porous space, Shale, Unconventional resources

References

- [1] Bushnev, D. A., Burdel'naya, N. S., & Zhuravlev, A. V. (2017). Organic matter in Upper Devonian deposits of the Chernyshev Ridge. *Geochemistry International*, 55(6), 548-558.
- [2] Kayukova, G. P., Kiyamova, A. M., Mikhailova, A. N., Kosachev, I. P., Petrov, S. M., Romanov, G. V., ... Vakhin, A. V. (2016). Generation of Hydrocarbons by Hydrothermal Transformation of Organic Matter of Domanik Rocks. *Chemistry and Technology of Fuels and Oils*, 52(2), 149-161.
- [3] Fadeeva, N. P., Kozlova, E. V., Poludetkina, E. N., Shardanova, T. A., Pronina, N. V., Stupakova, A. V., Khomyak, A. N. (2016). The hydrocarbon-generation potential of the Domanik rocks in the Volga-Ural petroliferous basin. *Moscow University Geology Bulletin*, 71(1), 41-49.
- [4] Gatovskii, Y. A., Stupakova, A. V., Kalmykov, G. A., Korobova, N. I., Suslova, A. A., Sautkin, R. S., & Kalmykov, D. G. (2016). New data on the biostratigraphy and facies types of upper Devonian Domanik sections in the Volga-Ural Basin. *Moscow University Geology Bulletin*, 71(1), 58-70.
- [5] Gilyazetdinova, D. R., & Korost, D. V. (2016). Transformation of the pore space during the simulation of the generation of hydrocarbon fluids in the Domanik horizon of the South-Tatar crest as an example. *Moscow University Geology Bulletin*, 71(1), 81- 88.

- [6] Girshgorn L. Sh., Zhamoyda A. I., Kovalevsky O. P., Oleinikov A. N., Prozorovskaya E. L., Hramov A. N., Shkatova, V. K. *Obschaya stratigraficheskaya shkala Rossii*. VSEGEI, Saint Petersburg, 2006.
- [7] International Chronostratigraphic Chart (2017). Published online by the IUGS International Commission on Stratigraphy. <http://www.stratigraphy.org/index.php/ics-chart-timescale>, 2017.
- [8] Kadyrov R.I., Platov B.V. (2018). Paleotectonic analysis of the sedimentary cover along the seismic profile "Tatseis" on the territory of the Republic of Tatarstan, Russia, *Proceedings of Kazan Golovkinsky Stratigraphic Meeting 2017*, 1, 349-355.
- [9] Bushnev, D. A., Burdel'naya, N. S., Ponomarenko, E. S., & Zubova Kiryukhina, T. A. (2016). Anoxia in the Domanik basin of the Timan--Pechora region. *Lithology and Mineral Resources*, 51(4), 283-289.